

Detail Procedures for Percutaneous coronary intervention

	Admission Day	First day of PCI		Second day of PCI
	Before Examination	Examination Day		The first day after examination
Tests	<ul style="list-style-type: none"> ■ EKG ■ Chest X-ray ■ Sign in information consent 	<ul style="list-style-type: none"> ■ Blood withdrawal 		<ul style="list-style-type: none"> ■ Administer medicine
Procedures	<ul style="list-style-type: none"> ■ Skin preparation ■ Injecting vein catheter and take blood to check-up. 	<ul style="list-style-type: none"> ■ Changing Operation clothes 	<ul style="list-style-type: none"> ■ The wound of groin or wrist compression to stop bleeding. 	<ul style="list-style-type: none"> ■ Dressing change
Medication		<ul style="list-style-type: none"> ■ intravenous fluid supply 		
Daily activity	<ul style="list-style-type: none"> ■ No limit action 	<ul style="list-style-type: none"> ■ No limit action before operation ■ Void before operation 	<ul style="list-style-type: none"> ■ Staying in the bed for 8 or 2 hours 	<ul style="list-style-type: none"> ■ Low sodium
Diet	<ul style="list-style-type: none"> ■ Low sodium ■ No food or soft drink after midnight. 	<ul style="list-style-type: none"> ■ Nothing per mouth ■ No meal or drink after breakfast for patient's scheduled at afternoon. 	<ul style="list-style-type: none"> ■ If there's no discomfort after operation, patient could start eating. 	<ul style="list-style-type: none"> ■ Low salt diet
Notification	<ul style="list-style-type: none"> ■ Explaining general notification <p>1. Inserting catheter from groin or wrist artery and follow the vessel to the heart after antiseptics and local anesthesia.</p> <p>2. Hotness sensation may follow after injection of contrast medicine. Don't be panic.</p> <p>Don't be nervous and you could take deep breath, the feeling of hot will disappear soon.</p> <p>3. You are awake during whole procedure. Please relax yourself with taking deep breath.</p> <p>4. Notify doctor if extreme coldness, high factor, SOB, or palpitation immediately.</p>	<ul style="list-style-type: none"> ■ Nurses will measure blood pressure, heart rate and healing of wound ■ People who is examined thru groin puncture : During sand-bag compression, keep in supine position and turn right by using wooden roll. ■ People who is examined thru radial artery puncture: Bed rest for 2 hours is enough. ■ During bed time, you need pee on the bed by using urine pot. You can't leave the bed for avoiding bleeding. ■ Inform nurse if you have any problem. 		<ul style="list-style-type: none"> ■ Guide-line for wound care: Don't wet the wound and take off the swab by yourself ■ To understand homecare notification after leaving hospital
Remarks	The list above are general regulations of hospitalization, the doctors will adjust according to individual actual condition.			